

PROSTATITIS

Patient Fact Sheet

Introduction

The prostate is a gland located at the base of the bladder, surrounding part of the urethra (the tube that carries urine from the bladder) in men. The prostate also functions in reproduction by producing part of the seminal fluid, which helps to transport sperm. Prostatitis is a common problem, affecting up to 25% of all men, but the diagnosis and treatment can be difficult. *Acute prostatitis* and *chronic bacterial prostatitis* are usually caused by the same types of bacteria that cause urinary tract infections. Prostatitis may also be related to sexually transmitted diseases like chlamydia, gonorrhea, or HIV. The most common type of prostatitis is *chronic prostatitis/chronic pelvic pain syndrome (CP/CPPS)*, which is most likely not caused by infection. CP/CPPS is recurrent, the exact cause is not known, and treatment can be challenging. The least common type of prostatitis is *asymptomatic inflammatory prostatitis*.

Acute prostatitis and Chronic bacterial prostatitis Symptoms

- Frequent need to urinate
- Feeling the need to urinate urgently
- Inability to urinate completely or dribbling urine
- Blood in urine or semen
- Low back pain
- Burning and/or pain when passing urine
- Rectal or scrotal area pain
- Groin pain
- Lower abdominal pain
- Fever and chills
- Flu-like symptoms
- Inability to get or keep an erection

Signs your health care provider may find on exam

- Urine studies reveal white blood cells, red blood cells, and/or bacteria
- Blood tests may show infection throughout the body in acute prostatitis (sepsis)
- Cultures may reveal STDs or other infection (chlamydia, gonorrhea, fungal infection)
- Cystoscopy (exam of the prostate by inserting a scope through the penis and urethra into the bladder) may reveal an enlarged prostate

- Pain with digital rectal exam (insertion of gloved finger into rectum to examine prostate)
- Discharge from the penis during prostate exam
- PSA (prostate specific antigen—blood test) may be elevated
- Fever (temperature greater than 101 degrees Fahrenheit)
- Tenderness in the lower abdomen and/or scrotum

Treatment

- Acute prostatitis sometimes requires hospitalization and intravenous (IV) antibiotics.
- Oral antibiotics are prescribed for both acute and chronic bacterial prostatitis—usually for at least 4-6 weeks, sometimes longer. It is important to take medication exactly as prescribed and to complete the medication to prevent the infection from recurring.
- Medication to relax the prostate may be given for symptoms of urgency/frequency.
- Medication to shrink the prostate size may be prescribed if prostate is enlarged.
- Anti-inflammatory medications like ibuprofen or naproxen may be prescribed for pain.
- If a man's prostate is very large (BPH) and the bacterial prostatitis keeps recurring, surgery to reduce the size of the prostate may be indicated.
- See your health care provider immediately if the above symptoms continue or reoccur

Chronic prostatitis/Chronic Pelvic Pain Syndrome (CP/CPPS) Symptoms

- Recurrent or ongoing pelvic pain, rectal pain, scrotal pain
- Low back pain
- Abdominal pain
- Blood in urine and/or semen
- Prostate area feels "swollen" or uncomfortable
- Pain with urination
- Painful intercourse
- Difficulty or hesitation with urination
- Depression
- CP/CPPS is often associated with other types of chronic pain, like chronic fatigue, fibromyalgia, and irritable bowel syndrome

This material is for educational purposes only and should in no way be taken to be the practice or provision of medical, nursing or professional healthcare advice or services. The information should not be used in place of a visit, call, consultation or advice of your physician, nurse or other health care provider. The information obtained herein is not exhaustive and does not cover all aspects of the specific disease, ailment, physical condition or their treatments. Should you have any health care related questions, please call or see your physician, nurse or other health care provider promptly.

The Society of Urologic Nurses and Associates is a professional organization committed to excellence in evidence-based clinical practice, research, and education of its members, patients, families, and the community.

© 2013 Society of Urologic Nurses and Associates



For more information, contact:

Society of Urologic Nurses and Associates

East Holly Avenue Box 56

Pitman, NJ 08071-0056

Phone 888-TAP-SUNA or 856-256-2335

suna@ajj.com

www.suna.org

Signs your health care provider may find on exam

- Mild prostate tenderness on digital rectal exam
- Muscle spasms on digital rectal exam
- Pelvic floor muscle problems
- Usually there is no fever, no chills
- No bacterial infection
- It can be very frustrating for CP/CPPS patients that often there are minimal or no significant abnormal findings on any of the tests or exams that are done.

Treatment possibilities—there is no definite, accepted treatment for CP/CPPS

- Antibiotics do not usually help CP/CPPS and use of antibiotics can lead to resistance (which means the antibiotics won't work when you need them)
- Anti-inflammatory medications like ibuprofen or naproxen can help relieve pain
- If the prostate is enlarged, medication that shrinks the prostate may help.
- Medication to relax the prostate may be used if urination is a problem.
- Physical therapy for pelvic floor muscle rehabilitation
- Myofascial trigger point release
- Prostate massage
- Sitz baths—sitting in a tub of warm water for 15-20 minutes
- Biofeedback
- Physical exercise
- Maintain an open line of communication with your clinician—tell your health care provider when your symptoms improve and also report if or when your symptoms worsen or recur. The same treatment does not work for everyone.

Asymptomatic inflammatory prostatitis

- There are no symptoms
- Elevated white blood cells are found in the patient's ejaculation fluid.
- This type of prostatitis is usually found when a man has a prostate biopsy or infertility or cancer workup.
- Asymptomatic inflammatory prostatitis is not usually treated.

Prevention of Prostatitis

- Often the exact cause for prostatitis cannot be isolated, but there are a few suggestions for prevention.
- Maintain good personal hygiene. Keep the genital area clean.
- Sexually transmitted diseases can cause prostatitis. Engage in safe sexual practices—use barrier protection/condoms.
- Seek treatment promptly if you have signs or symptoms of a urinary tract infection.
- There is some evidence suggesting that a diet rich in fruits, vegetables and fiber may improve prostate health.